

Advances on Asymptotic Stability of Impulsive Stochastic Evolution Equations

Sheila A. Bishop, Kanayo S. Eke, Hilary I. Okagbue

Department of Mathematics
Covenant University
Ota, Ogun State, Nigeria

email: sheila.bishop@covenantuniversity.edu.ng, sheilabishop95@yahoo.com

(Received April 18, 2020, Revised May 18, 2020, Accepted May 26, 2020)

Abstract

Asymptotic stability of impulsive stochastic evolution equations (ISEEs) with nonlocal initial conditions is studied. Stability analysis of impulsive stochastic differential equations with local initial conditions has received substantial attention. While relatively, little has been done on asymptotic stability with nonlocal initial conditions. Considering the advantage nonlocal initial value problems have over local initial value problems, we study the asymptotic stability of ISEEs with nonlocal initial conditions. Using the contraction mapping technique and by perturbing the local initial condition, asymptotic stability of ISEEs is established.

1 Introduction

Recently Impulsive evolution differential equations (stochastic, ordinary, partial, functional, quantum, etc.) with local and nonlocal initial conditions have become more relevant in so many mathematical models [1-23] of real processes and phenomena studied in virtually all branches of science, technology, social sciences, etc. There has been an appreciable development in impulse theory, particularly concerning impulsive differential equations with

Key words and phrases: Asymptotic stability, Evolution operator, Stochastic differential equation, Fixed point, Nonlocal initial condition, Mild solution, Impulse effects.

AMS (MOS) Subject Classifications: 37L05, 47J35, 47H10, 93D20.
ISSN 1814-0432, 2021, <http://ijmcs.future-in-tech.net>

fixed moments [4-5, 9, 11, 14-15, 17-18, 20, 24-29, 32]. The study of the theory of stability of stochastic differential equations (SDEs) has been investigated by the Lyapunov functional method, energy estimates method, etc. (see [22, 30] and the references therein). The Lyapunov function has been used as a vital instrument to study the qualitative properties of the trivial solutions of differential equations [5, 11, 13, 15, 17]. Oleksiy [13] considered the asymptotic stability of a system of SDEs which has a zero solution by assuming the existence of a positive definite function. Wu et al. [4] introduced a time-varying discontinuous Lyapunov-Krasovskii functional to establish the stability of an impulsive system. The Lyapunov method, though extensively used, has some difficulties especially when studying asymptotic stability of SDEs (see [22] and the references therein). By applying some novel techniques, such as the fixed point approach, some of these difficulties have been addressed. Jing et al. [8] worked on the existence and exponential stability for a class of neutral stochastic partial differential equations using the Banach fixed point principle. Recently, Sufang et al. [9] investigated the existence and exponential stability of mild solutions for a class of impulsive stochastic differential equations (ISDEs) driven by a functional Brownian motion via Monch fixed point theorem. By using the fixed point approach Bishop et al. [10] discussed Ulam's type of stability of quantum stochastic differential equations (QSDEs) which led to some interesting results. Evolution differential equations have found practical applications in virtually all fields of science and technology [18, 21, 23, 31]. Also, problems with nonlocal initial conditions have been shown to have an advantage over traditional initial value problems ([9, 13, 29]). Motivated by the results in the References [6, 8, 9, 22, 23], we study the stability of solutions of ISEEs with nonlocal initial conditions. Hence, the results in this paper would generalize results on asymptotic stability of stochastic differential equations with initial local conditions and will have application to a wider class of equations. The rest of this paper consists of two sections: section 2 has to do with preliminary results, while the major results are discussed in section 3.

2 Preliminaries

Let (Ω, Σ, P) be a complete probability space, $\{\Sigma_t\}_{t \geq 0}$ a normal filtration. H, Y are separable Hilbert spaces with the usual norms $\|\cdot\|_H$ and $\|\cdot\|_Y$ respectively. Let Q be a Wiener process on (Ω, Σ, P) with $\text{tr}Q < \infty$. Let $L(H, Y)$ denote the space of linear operators that are bounded, $C(H, H)$ the space of

all continuous functions defined on H , $\Sigma_t = \Sigma_t^w$ a sigma algebra and L_2^0 is as defined in [22].

Of interest here is the following impulsive stochastic evolution equations (ISEEs):

$$\begin{aligned}
 d\phi(t) &= [A\phi(t) + F(t, \phi(t))]dt + G(t, \phi(t))dw(t) \\
 t \in I &= [0, T] \subseteq \mathbb{R}_+, t \neq t_k \\
 \Delta\phi(t_k) &= J_k(\phi(t_k)), t = t_k, k = 1, \dots, m \\
 \phi(0) &= \phi_0 + g(\phi), t \in [0, T], 0 \leq T < \infty.
 \end{aligned}
 \tag{2.1}$$

Here $J_k \in C(H, H)$, A is defined as the infinitesimal generator of a semi-group of bounded linear operators $S(t), t \geq 0$, in H . The function $g : C([0, T], H) \rightarrow H$ is continuous and compact such that $E \{sup \|\phi(t)\|_H\} < +\infty$.

$\phi(t_k^+), \phi(t_k^-)$ are the right and left limits of $\phi(t)$ at t_k , where $0 < t_1 < t_2 < \dots < t_k < T$. $\Delta\phi(t_k) = \phi(t_k^+) - \phi(t_k^-)$ is the jump in the state ϕ at t_k . The maps $F \in (\mathbb{R}_+ \times H, H)$ and $G \in (\mathbb{R}_+ \times H, L(H, Y))$ are Borel measurable.

Definition 2.1. A stochastic process $\phi(t)$ is called a mild solution of (2.1) if:

1. $\phi(t)$ is Σ_t - adapted,
2. $\phi(t) \in H$ admits jumps for $t \in [0, T]$ and
3. $\phi(t)$ satisfies the integral equation

$$\begin{aligned}
 \phi(t) &= S(t)[\phi(0) + g(\phi)] \\
 &+ \int_0^t S(t-s) (F(s, \phi(s))ds + G(s, \phi(s))dw(s)) \\
 &+ \sum_{0 < t_k < t} S(t-t_k)J_k(\phi(t_k)).
 \end{aligned}
 \tag{2.2}$$

Definition 2.2. Equation (2.2) is stable in the p th moment if for any arbitrary $\epsilon > 0$ we can find a $\delta > 0$ such that $\|\phi_0\|_H < \delta$ and $E \{sup_{t \geq 0} \|\phi(t)\|_H^p\} < \epsilon, p \geq 2 \in \mathbb{N}$.

Definition 2.3. Equation (2.2) is asymptotically stable in the p th moment if Definition 2.2 holds such that for any $\phi_0 \in H$,

$$\lim_{t \rightarrow \infty} E \left\{ \sup_{t \in [0, T]} \|\phi(t)\|_H^p \right\} = 0.$$

We state the following well known results.

Lemma 2.4. For any $\gamma \geq 1$ and for arbitrary L_2^0 - valued process $\vartheta(\cdot)$,

$$\sup_{s \in [0, t]} E \left\| \int_0^s \vartheta(u) dw(u) \right\|_H^{2\gamma} \leq (\gamma(2\gamma - 1))^\gamma \left(\int_0^t (E \|\vartheta(s)\|_{L_2^0}^{2\gamma})^{1/\gamma} ds \right)^\gamma.$$

Theorem 2.5. Let Ψ be a contraction operator on a Banach space B , then there exists a unique point $x \in B$ for which $\Psi(x) = x$.

Remark 2.6. The proof of these results can be found in [22] and the references therein.

3 Major Results

In this section, we make some assumptions on the nonlocal conditions and establish asymptotic stability of solutions of the given problem by transforming the problem into a fixed point problem. We show that the problem has a fixed point in the state space using the contraction mapping method and we are done.

We assume the following conditions:

1. $F(t, 0) = 0$, $G(t, 0) = 0$ and $J_k(0) = 0$, $k = 1, 2, \dots, m$.
2. The zero solution of (2.1) exists when $\phi = 0$ and $g(0) = 0$.
3. The map $\psi(t, w) : [0, \infty) \times \Omega \rightarrow \mathbb{R}$ is continuous in t such that $E \|\psi(t, w)\|_H^p \rightarrow 0$ as $t \rightarrow \infty$.

Next, we state the following major conditions:

Let $c_k > 0$, $M \geq 0$, $L_g \geq 0$, K and $L > 0$ (are Lipschitz constants) such that

- (S₁) for each $\phi, \varphi \in H$, $\|J_k(\phi) - J_k(\varphi)\| \leq c_k \|\phi - \varphi\|$, $k = 1, 2, \dots, m$;
- (S₂) $\|S(t)\|_H \leq M e^{-at}$, $t \geq 0$, $a > 0$;
- (S₃) $\|g(\phi) - g(\varphi)\|_H \leq L_g \|\phi - \varphi\|_H$;
- (S₄) $\|F(t, \phi) - F(t, \varphi)\|_H \leq K \|\phi - \varphi\|_H$, $\|G(t, \phi) - G(t, \varphi)\|_H \leq L \|\phi - \varphi\|_H$;
- (S₅) $4^{p-1} M^p (L_g a^{-p} + K^p a^{-p} + L^p l_p (2a)^{-p/2} + L_k) < 1$,

where $l_p = (p(p-1)/2)^{p/2}$ and $L_k = e^{-apT} E (\sum_{k=1}^m \|c_k\|_H)$;

and

- (S₆) for any arbitrary $\epsilon > 0$ and $\phi(t) \in H$, $t \in [0, T]$, $\|\phi(t)\|_H^p < \epsilon$.

Theorem 3.1. Assume that the conditions (S₁) – (S₆) are satisfied. Then the mild solution of the nonlocal problem (2.1) is asymptotically stable in the p th moment.

Proof. Let the operator $\Gamma : C(I, B) \rightarrow C(I, B)$. We define a mapping Γ on

$C(I, B)$ by

$$\begin{aligned} \Gamma(\phi)(t) &= S(t)[\phi_0 + g(\phi)] + \int_0^t S(t-s)F(s, \phi(s))ds \\ &\quad + \int_0^t S(t-s)G(s, \phi(s))ds + \sum_{0 < t_k < t} S(t-t_k)J_k(\phi(t_k)) \\ &= \sum_{i=1}^4 \Pi_i(t), t \geq 0, i = 1, 2, 3, 4. \end{aligned} \tag{3.1}$$

It suffices to show that Γ is a contraction operator on B (see [22] and the references therein). We show this in the following steps.

Step 1: The operator Γ is continuous.

Let $0 < \tau < \epsilon$. Then

$$E \|\Gamma(\phi)(t_1 + \tau) - \Gamma(\phi)(t_1)\|_H^p \leq 4^{p-1} \sum_{i=1}^4 E \|\Pi_i(t_1 + \tau) - \Pi_i(t_1)\|_H^p.$$

and

$$E \|\Pi_i(t_1 + \tau) - \Pi_i(t_1)\|_H^p \rightarrow 0$$

as $\tau \rightarrow 0, i = 1, 2, 3, 4$.

Using the well known Holder's inequality and Lemma 2.4, we obtain

$$\begin{aligned} E \|\Pi_3(t_1 + \tau) - \Pi_3(t_1)\|_H^p &\leq 3^{p-1} l_p [E \|((S(t_1 + \tau) - S(t_1))g(\phi))\|_H^p]^{(2/p)} \\ &\quad + 3^{p-1} l_p \left[\int_0^{t_1} (E \|(S(t_1 + \tau - s) - S(t_1 - s))G(s, \phi(s))\|_H^p)^{(2/p)} ds \right]^{(p/2)} \\ &\quad + 3^{p-1} l_p \left[\int_{t_1}^{t_1 + \tau} (E \|S(t_1 + \tau - s)G(s, \phi(s))\|_H^p)^{(2/p)} ds \right]^{(p/2)} \rightarrow 0 \end{aligned} \tag{3.2}$$

as $\tau \rightarrow 0$.

Showing that the operator Γ is continuous on $[0, T]$.

Step 2: $\Gamma(B) \subset B$.

Using (3.1), gives

$$\begin{aligned} E \|(\Gamma\phi)(t)\|_H^p &\leq 4^{p-1} E \|S(t)[\phi_0 + g(\phi)]\|_H^p \\ &\quad + 4^{p-1} E \left\| \int_0^t S(t-s)F(s, \phi(s))ds \right\|_H^p \\ &\quad + 4^{p-1} E \left\| \int_0^t S(t-s)G(s, \phi(s))dw_s \right\|_H^p \\ &\quad + 4^{p-1} \sum_{0 < t_k < t} E \|S(t-t_k)J_k(\phi(t_k))\|_H^p. \end{aligned} \tag{3.3}$$

Using (S_1) - (S_3) yields

$$4^{p-1}E \|S(t)[\phi_0 + g(\phi)]\|_H^p \leq M^p e^{-pat} \|\phi_0 + g(\phi)\|_H^p \rightarrow 0 \quad (3.4)$$

and

$$4^{p-1} \sum_{0 < t_k < t} E \|S(t - t_k)J_k(\phi(t_k))\|_H^p \leq M^p e^{-pat} \|J_k(\phi(t_k))\|_H^p \rightarrow 0 \quad (3.5)$$

as $t \rightarrow \infty$.

By using (S_1) , (S_4) and Holder's inequality, we derive

$$\begin{aligned} 4^{p-1}E \left\| \int_0^t S(t-s)F(s, \phi(s))ds \right\|_H^p &\leq 4^{p-1}M^p K^p \left[\int_0^t e^{-a(t-s)} ds \right]^{p-1} \\ &\quad \times \int_0^t e^{-a(t-s)} E \|\phi(s)\|_H^p ds \\ &\leq 4^{p-1}M^p K^p a^{1-p} \int_0^t e^{-a(t-s)} E \|\phi(s)\|_H^p ds. \end{aligned} \quad (3.6)$$

Now by (S_6) and (3.4) we get

$$\begin{aligned} 4^{p-1}E \left\| \int_0^t S(t-s)F(s, \phi(s))ds \right\|_H^p &\leq 4^{p-1}M^p K^p a^{1-p} e^{-at} \int_0^{t_1} e^{as} E \|\phi(s)\|_H^p ds \\ &\quad + 4^{p-1}M^p K^p a^{-p} \epsilon. \end{aligned} \quad (3.7)$$

By the hypothesis of Theorem 3.1 as $t \rightarrow \infty$, $e^{-at} \rightarrow 0$. Now, for $0 < t_1 < t$, we obtain

$$4^{p-1}M^p K^p a^{1-p} e^{-at} \int_0^{t_1} e^{as} E \|\phi(s)\|_H^p ds \leq \epsilon - 4^{p-1}M^p K^p a^{-p} \epsilon. \quad (3.8)$$

Hence, using (3.7) and (3.8), yields

$$4^{p-1}E \left\| \int_0^t S(t-s)F(s, \phi(s))ds \right\|_H^p < \epsilon.$$

Which implies that

$$4^{p-1}E \left\| \int_0^t S(t-s)F(s, \phi(s))ds \right\|_H^p \rightarrow 0 \text{ as } t \rightarrow \infty. \quad (3.9)$$

Similarly, for $\phi \in B$ and $t \in [0, T]$,

$$4^{p-1}E \left\| \int_0^t S(t-s)G(s, \phi(s))dw(s) \right\|_H^p \leq 4^{p-1}l_p M^p L^p \times \left[\int_0^t e^{-2a(t-s)} (E \|\phi(s)\|_H^p)^{2/p} ds \right]^{p/2}. \tag{3.10}$$

So that in the same vain as (3.9) and (3.10), we get

$$4^{p-1}E \left\| \int_0^t S(t-s)G(s, \phi(s))dw(s) \right\|_H^p \rightarrow 0 \text{ as } t \rightarrow \infty. \tag{3.11}$$

Therefore, substituting (3.4), (3.5), (3.9) and (3.11) in (3.3), we see that $E \|\Gamma\phi(t)\|_H^p \rightarrow 0$ as $t \rightarrow \infty$. Hence, $\Gamma(B) \subset B$.

Step 3: The operator Γ is a contraction mapping on B .

For $\phi, \varphi \in B, t \in [0, T]$,

$$\begin{aligned} \sup_{s \in [0, T]} E \|\Gamma\phi(t) - \Gamma\varphi(t)\|_H^p &\leq 4^{p-1}E \|S(t)[g(\phi) - g(\varphi)]\|_H^p \\ &+ 4^{p-1} \sup_{s \in [0, T]} E \left\| \int_0^t S(t-s)[F(s, \phi(t)) - F(s, \varphi(s))]ds \right\|_H^p \\ &+ 4^{p-1} \sup_{s \in [0, T]} E \left\| \int_0^t S(t-s)[G(s, \phi(t)) - G(s, \varphi(s))]dw(s) \right\|_H^p \\ &+ 4^{p-1} \sup_{s \in [0, T]} E \left\| \sum_{0 < t_k < t} S(t-t_k)[J_k(\phi(t_k)) - J_k(\varphi(t_k))] \right\|_H^p \\ &\leq [4^{p-1}M^p(L_g a^{-p} + K^p a^{-p} + L^p l_p (2a)^{-p/2} + L_k)] \\ &\times \left(\sup_{t \in [0, T]} E \|\phi(t) - \varphi(t)\|_H^p \right). \end{aligned} \tag{3.12}$$

Therefore,

$$\sup_{s \in [0, T]} E \|\Gamma\phi(t) - \Gamma\varphi(t)\|_H^p \leq \sup_{t \in [0, T]} E \|\phi(t) - \varphi(t)\|_H^p,$$

where $4^{p-1}M^p(L_g a^{-p} + K^p a^{-p} + L^p l_p (2a)^{-p/2} + L_k) < 1$ showing that Γ is a contraction mapping. This implies that a fixed point $\phi(\cdot) \in B$ which is a solution of the nonlocal problem (2.1) exists and $E \|\phi(t)\|_H^p \rightarrow 0$ as $t \rightarrow \infty$.

The proof is therefore complete and by Definition 2.3 we conclude that the solution of the nonlocal problem (2.1) is asymptotically stable.

Acknowledgment. The authors will like to thank the anonymous reviewers for their valuable comments.

References

- [1] S. A. Bishop, M. O. Ogundiran, O. P. Ogundile, On Stability of Quantum Stochastic Differential Equation, *Int. Journal Mech. Engg. Tech.*, **9**, no. 9, (2018), 367–374.
- [2] S. A. Bishop, E. O. Ayoola, G. J. Oghonyon, Existence of mild solution of impulsive quantum stochastic differential equation with nonlocal conditions, *Anal. Math. Phys.*, **7**, no. 3,(2017), 255–265.
- [3] T. Caraballo, M. J. Garrido-Atienza, T. Taniguchi, The existence and exponential behavior of solutions to stochastic delay evolution equations with a fractional Brownian motion, *Nonlinear Analysis: Theory, Methods and Applications*, **74**, no. 11, (2011), 3671–3684.
- [4] W. H. Chen, W. X. Zheng, Input-to-state stability for networked control systems via an improved impulsive system approach, *Automatica*, **47**, no. 4, (2011), 789–796.
- [5] P. Cheng, F. Deng, Fengqi Yao, Almost sure exponential stability and stochastic stabilization of stochastic differential systems with impulsive effects, *Nonlinear Analysis: Hybrid Systems*, **30**, (2018), 106–117.
- [6] P. Chen, A. Abdelmonem, Y. Li, Global existence and asymptotic stability of mild solutions for stochastic evolution equations with nonlocal initial conditions, *J. Integral Equations and Applications*, **29**, no. 2, (2017), 325–348.
- [7] J Cui, L. Yan, Successive approximation of neutral stochastic evolution equations with infinite delay and Poisson jumps, *Applied Mathematics and Computation*, **218**, no. 12, 20120, 6776–6784.
- [8] J. Cui, L. Yan, X. Sun, Exponential stability for neutral stochastic partial differential equations with delays and Poisson jumps, *Statistics and Probability Letters*, **81**, no. 12, (2011), 1970–1977.

- [9] S. Deng, X. B. Shu, J. Mao, Existence and exponential stability for impulsive neutral stochastic functional differential equations driven by fBm with noncompact semigroup via Mönch fixed point, *Journal of Mathematical Analysis and Applications*, **467**, no. 1, (2018), 398–420.
- [10] S. Diaz-Infante, S. Jerez, n Convergence and asymptotic stability of the explicit Steklov method for stochastic differential equations, *Journal of Computational and Applied Mathematics*, **291**, (2016), 36–47.
- [11] L. Gao, D. Wang, G. Zong, Exponential stability for generalized stochastic impulsive functional differential equations with delayed impulses and Markovian switching, *Nonlinear Analysis: Hybrid Systems*, **30**, (2018), 199–212.
- [12] E. Hernandez, Existence results for partial neutral functional differential equations with nonlocal conditions, *Dynam. Systems Appl.*, **11**, no. 2, (2002), 241–252.
- [13] O. Ignatyev, Partial asymptotic stability in probability of stochastic differential equations, *Statistics and Probability Letters*, **79**, no. 5, (2009), 597–601.
- [14] J. Liang, H.L. James, Ti-Jun Xiao, Nonlocal impulsive problems for nonlinear differential equations in Banach spaces, *Mathematical and Computer Modelling*, **49**, (2009), 798–804.
- [15] B. Li, Stability of stochastic functional differential equations with impulses by an average approach *Nonlinear Analysis: Hybrid Systems*, **29**, (2018), 221–233.
- [16] L. Liu, F. Deng, pth moment exponential stability of highly nonlinear neutral pantograph stochastic differential equations driven by Lévy noise, *Applied Mathematics Letters*, **86**, (2018), 313–319.
- [17] S. Liu, F. Chen, Z. Wang, Existence of global L_p solutions to a symmetric system of Keyfitz–Kranzer type, *Applied Mathematics Letters*, **52**, (2016), 96–101.
- [18] M. McKibben, *Discovering Evolution Equations with Applications: Volume 1, Deterministic Equations* (Chapman and Hall/CRC Applied Mathematics and Nonlinear Science), 1st Edition, 2010, London, NY.

- [19] A. Paicu, I.I. Vrabie, A class of nonlinear evolution equations subjected to nonlocal initial conditions, *Nonlinear Analysis: Theory, Methods and Applications*, **72**, no. 11, (2010), 4091–4100.
- [20] B. Pei, Y. Xu, Mild solutions of local non-Lipschitz stochastic evolution equations with jumps, *Applied Mathematics Letters*, **52**, (2016), 80–86.
- [21] B. L. Rozovskii, *Stochastic Differential Equations: Theory and Applications*, Interdisciplinary Mathematical Sciences, A Volume in Honor of Professor Boris L Rozovskii, April 2007, 420pp.
- [22] R. Sakthivel, J. Luo, Asymptotic stability of nonlinear stochastic differential equations, *Statistics and Probability Letters*, **79**, (2009), 1219–1223.
- [23] M. C. Veraar, Non-autonomous stochastic evolution equations and applications to stochastic partial differential equations, *J. Evol. Eq.*, **10**, (2010), 85–127.
- [24] F. Wu, G. Yin, L.Y. Wang, Moment exponential stability of random delay systems with two-time-scale Markovian switching, *Nonlinear Analysis: Real World Applications*, **13**, no. 6, (2012), 2476–2490.
- [25] X. Wu, W. Zhang, Y. Tang, p th Moment stability of impulsive stochastic delay differential systems with Markovian switching, *Communications in Nonlinear Science and Numerical Simulation*, **18**, no. 7, (2013), 1870–1879.
- [26] L. Xu, J. Luo, Viability for stochastic functional differential equations in Hilbert spaces driven by fractional Brownian motion *Applied Mathematics and Computation*, **341**, (2019), 93–110.
- [27] L. Xu, Z. Li, Stochastic fractional evolution equations with fractional brownian motion and infinite delay, *Applied Mathematics and Computation*, **336**, (2018), 36–46.
- [28] L. Xu, Z. Dai, D. He, Exponential ultimate boundedness of impulsive stochastic delay differential equations, *Applied Mathematics Letters*, **85**, (2018), 70–76.
- [29] H. Yang, M. Song, M. Liu, Strong convergence and exponential stability of stochastic differential equations with piecewise continuous arguments

- for non-globally Lipschitz continuous coefficients, *Applied Mathematics and Computation*, **341**, (2019), 111–127.
- [30] T. Zhang, H. Chen, C. Yuan, T. Caraballo, On the asymptotic behavior of highly nonlinear hybrid stochastic delay differential equations, *Dynamical Systems - B*, **24**, no. 10, (2019), 5355–5375.
- [31] S. Zheng, *Nonlinear Evolution Equations, Monographs and Surveys in Pure and Applied Mathematics*, Chapman and Hall/CRC, London, NY, (2004), 304pp.
- [32] Y. Zhou, F. Jiao, Nonlocal Cauchy problem for fractional evolution equations, *Nonlinear Analysis: Real World Applications*, **11**, no. 5, (2010), 4465–4475.